

Name

Class

# MATHS TEACHER HUB

www.MathsTeacherHub.com

## Bounds

(9 – 1) Topic booklet

## HIGHER

These questions have been collated from previous years GCSE Mathematics papers.

**You must have:** Ruler graduated in centimetres and millimetres, protractor, pair of compasses, pen, HB pencil, eraser.

Total Marks

### Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- Diagrams are NOT accurately drawn, unless otherwise indicated.
- You must **show all your working out.**
- If the question is a **1F** question you are not allowed to use a calculator.
- If the question is a **2F** or a **3F** question, you may use a calculator to help you answer.

### Information

- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

**Answer ALL questions**

**Write your answers in the space provided.**

**You must write down all the stages in your working.**

16  $p = \sqrt{\frac{2e}{f}}$

$e = 6.8$  correct to 1 decimal place.

$f = 0.05$  correct to 1 significant figure.

Work out the upper bound for the value of  $p$ .

Give your answer correct to 3 significant figures.

You must show all your working.

**16** The petrol consumption of a car, in litres per 100 kilometres, is given by the formula

$$\text{Petrol consumption} = \frac{100 \times \text{Number of litres of petrol used}}{\text{Number of kilometres travelled}}$$

Nathan's car travelled 148 kilometres, correct to 3 significant figures.

The car used 11.8 litres of petrol, correct to 3 significant figures.


Nathan says,

“My car used less than 8 litres of petrol per 100 kilometres.”

Could Nathan be wrong?

You must show how you get your answer.

17



$a$  is 8.3 cm correct to the nearest mm  
 $b$  is 6.1 cm correct to the nearest mm


Calculate the upper bound for  $c$ .  
You must show your working.

..... cm

Specimen 2 – Paper 2H

**(Total for Question 17 is 4 marks)**

18 The diagram shows a cube.



$AH = 11.3$  cm correct to the nearest mm.

Calculate the lower bound for the length of an edge of the cube.  
You must show all your working.

..... cm

**18** A high speed train travels a distance of 487 km in 3 hours.

The distance is measured correct to the nearest kilometre.

The time is measured correct to the nearest minute.

By considering bounds, work out the average speed, in km/minute, of the train to a suitable degree of accuracy.

You must show all your working and give a reason for your answer.

.....km/minute

**18**  $m = \frac{\sqrt{s}}{t}$   $s = 3.47$  correct to 3 significant figures  
 $t = 8.132$  correct to 4 significant figures

By considering bounds, work out the value of  $m$  to a suitable degree of accuracy.  
Give a reason for your answer.

19  $D = \frac{u^2}{2a}$

$u = 26.2$  correct to 3 significant figures

$a = 4.3$  correct to 2 significant figures

- (a) Calculate the upper bound for the value of  $D$ .  
Give your answer correct to 6 significant figures.  
You must show all your working.

.....  
(3)

The lower bound for the value of  $D$  is 78.6003 correct to 6 significant figures.

- (b) By considering bounds, write down the value of  $D$  to a suitable degree of accuracy.  
You must give a reason for your answer.

.....  
.....  
(2)



**20**  $d = \frac{1}{8}c^3$

$c = 10.9$  correct to 3 significant figures.

By considering bounds, work out the value of  $d$  to a suitable degree of accuracy.  
Give a reason for your answer.

**21** The time period,  $T$  seconds, of a simple pendulum of length  $l$  cm is given by the formula

$$T = 2\pi \sqrt{\frac{l}{g}}$$

Katie uses a simple pendulum in an experiment to find an estimate for the value of  $g$ .

Here are her results.

$l = 52.0$  correct to 3 significant figures.

$T = 1.45$  correct to 3 significant figures.

Work out the upper bound and the lower bound for the value of  $g$ .

Use  $\pi = 3.142$

You must show all your working.

upper bound = .....

lower bound = .....

**21** Jackson is trying to find the density, in  $\text{g/cm}^3$ , of a block of wood.  
The block of wood is in the shape of a cuboid.

He measures

the length as 13.2 cm, correct to the nearest mm

the width as 16.0 cm, correct to the nearest mm

the height as 21.7 cm, correct to the nearest mm

He measures the mass as 1970 g, correct to the nearest 5 g.

By considering bounds, work out the density of the wood.

Give your answer to a suitable degree of accuracy.

You must show all your working and give a reason for your final answer.

22  $D = \frac{x}{y}$

$x = 99.7$  correct to 1 decimal place.

$y = 67$  correct to 2 significant figures.

Work out an upper bound for  $D$ .

**23** A race is measured to have a distance of 10.6 km, correct to the nearest 0.1 km.  
Sam runs the race in a time of 31 minutes 48 seconds, correct to the nearest second.

Sam's average speed in this race is  $V$  km/hour.

By considering bounds, calculate the value of  $V$  to a suitable degree of accuracy.  
You must show all your working and give a reason for your answer.